

MYCOTAXON

<http://dx.doi.org/10.5248/126.15>

Volume 126, pp. 15–22

October–December 2013

Hyphomycetes: *Linodochium sinense* sp. nov. and new records from fallen *Sycopsis sinensis* leaves in ChinaDE-WEI LI^{*1}, JINGYUAN CHEN² & YIXUN WANG²¹*The Connecticut Agricultural Experiment Station, Valley Laboratory,
153 Cook Hill Road, Windsor, CT 06095, USA*²*Institute of Forest Disease and Insect Control, Hubei Academy of Forestry,
1 Lion Peak, Jiufeng, Wuhan, Hubei, 430075, China** CORRESPONDENCE TO: dewei.li@ct.gov

ABSTRACT — Among hyphomycetes collected from dead leaves of *Sycopsis sinensis* in Houhe National Nature Reserve, Wufeng, Hubei, China, a sporodochial fungus new to science, *Linodochium sinense*, is described and illustrated. It develops yellow conidiomata and 2-celled conidia that are filiform, colorless, smooth, $34\text{--}40 \times 1.1\text{--}1.5\ \mu\text{m}$. *Candelabrum microsporum*, *Chalara affinis*, *Cylindrocladium madagascariense*, and *Subulispora britannica* are new records from China, while *Menisporopsis novae-zelandiae* is newly reported from mainland China.

KEY WORDS — anamorphic fungi, saprobe

Introduction

Houhe National Nature Reserve in Wufeng Tujia Autonomous County, Hubei, China is exceptional for its rich biodiversity and rare species. With over 2292 species of vascular plants including 29 endangered species (Song & Liu 1999, Li et al. 2005), the Reserve also possesses some species native only to China (Dang et al. 2004, Li et al. 2005). Its fungal diversity is almost entirely unexplored, and only two studies (on ascomycetes and polypores) have been conducted there (Li et al. 2007, Zhuang et al. 2007).

Sycopsis sinensis Oliv. (*Hamamelidaceae*) is a native evergreen broadleaf woody plant that grows sporadically in mixed evergreen broadleaf forests or bushes at elevations of 1000–2000 m in southern China (Li et al. 2003). The largest population of *S. sinensis* as dominant species has been reported in Houhe National Nature Reserve (Liu et al. 1999).

Two field trips were made to the Reserve to collect hyphomycetes in August 2008 and May 2009. Hyphomycetes on dead leaves of *S. sinensis* were studied

and categorized. Three new hyphomycetes from the two surveys have already been published, one of them from *S. sinensis* (Li et al. 2010, Li et al. 2011). Five additional hyphomycetes from *S. sinensis* are reported here, one as a new species, three as new records from China, and one as a new record from mainland China.

Materials & methods

Sporodochia, conidiophores and conidia of the fungus were mounted in lacto-fuchsin (0.1 g acid fuchsin, 100 ml 85% lactic acid; Carmichael 1955) or 85% lactic acid. Microscopic observations were made using bright field and Nomarski differential interference contrast optics. Photomicrographs were taken with an Olympus MicroFire digital camera (Goleta, CA). Fungal structures were measured with PictureFrame 2.3 software (Goleta, CA). Measurements of the fungal structures were statistically analyzed with Microsoft Office Excel 2010 with 95% confidence interval of means. The results are presented as ranges and mean $\pm$ standard deviation. Q = length/width ratio and n = the number of fungal structures measured. Herbarium acronyms follow Index Herbariorum (Thiers 2013).

Taxonomy

Linodochium sinense D.W. Li, Jing Y. Chen & Yi X. Wang, **sp. nov.**

FIGS 1–6

MYCOBANK MB 804385

Differs from other *Linodochium* spp. by conidiogenous cells that bear 1–3 (mostly 2) 1-septate conidia with dichotomous denticulate apices and conidial anastomoses that form x-shaped derivative conidia.

TYPE: China, Hubei, Wufeng, Houhe National Nature Reserve, on dead leaves of *Sycopsis sinensis*, 20 May 2009, Yixun Wang (**Holotype**: BPI 892531).

ETYMOLOGY: referring to China for its origin.

CONIDIOMATA sporodochial, circular to irregular, bright yellow when young, orange at maturity, scattered, hypophyllous, occasionally amphigenous, 200–300 μm diam. and 90–110 μm high. CONIDIOPHORES differentiated, erect, branched, flexuous or straight, colorless to pale yellow, smooth, septate, packed together. CONIDIOGENOUS CELLS integrated, terminal, holoblastic, straight or flexuous, bacilliform or ampulliform, sympodially elongated, smooth, thin-walled, colorless, $(11\text{--})12\text{--}16(\text{--}19) \times (1\text{--})1.2\text{--}1.4(\text{--}1.7) \mu\text{m}$ (mean = $14.4 \pm 1.8 \times 1.3 \pm 0.1$, $n = 30$), bearing mostly $(1\text{--})2(\text{--}3)$ conidia; apices of some conidiogenous cells ending with two flat-topped denticles of different lengths

FIGURES 1–8. *Linodochium sinense* (Holotype: BPI 892531). 1. Sporodochia on lower epidermis of a leaf of *Sycopsis sinensis*, 2. Cross-sections of sporodochia. 3. Conidiogenous cells and conidiophores. 4. Conidia on conidiogenous cells and four pairs of conidia on the right anastomosed into x-shaped derivative conidia. 5. An x-shaped derivative conidium. 6–7. Conidia on a conidiogenous cell with a dichotomous denticulate apex. 8. Three conidia on a conidiogenous cell with a trichotomous denticulate apex. Scale bars: 1 = 1 mm; 2 = 10 μm ; 3–8 = 5 μm .



forming a dichotomy, some with one denticle and one flat conidiogenous locus on the other side of the conidiogenous cell apex, rarely, conidiogenous cells ending in a trichotomous denticulate apex. CONIDIA 1-septate, filiform, colorless, smooth, guttulate, $(32-34-40(-45) \times (1-)1.1-1.5(-1.6) \mu\text{m}$ (mean = $37.5 \pm 3 \times 1.3 \pm 0.2$, $n = 30$), $Q = (22.5-)24.5-32.1(-39.3)$ (mean = 28.3 ± 3.8 , $n = 30$); some conidial pairs anastomosing to form x-shaped derivative conidia.

TELEOMORPH: unknown.

GEOGRAPHICAL DISTRIBUTION: Hubei, China.

HABITAT: dead leaves of *Sycopsis sinensis*.

NOTES: *Linodochium* was erected by Höhnelt (1909) and typified by *Linodochium hyalinum* (Lib.) Höhn. It remained monotypic for seven decades. Dyko & Sutton (1979) redescribed and illustrated the genus and *L. hyalinum*, its only species. Two years later, Minter & Holubová-Jechová (1981) added a second species, *L. formosum* Minter & Hol.-Jech. During the 1990's two more species were described: *Linodochium album* R.F. Castañeda & W.B. Kendr. (Castañeda & Kendrick 1990) and *L. hyalinum*, *L. sideroxyli* Dulym. et al. (Dulymamode et al. 1999). *Linodochium sinense* is the fifth species proposed in the genus. Much longer conidia differentiate *L. formosum* (80–140 μm with 2–4 septa; Dyko & Sutton 1979) and *L. hyalinum* (36–89 μm with <12 septa; Minter & Holubová-Jechová 1981) from *L. sinense*. White sporodochia and falcate conidia of *L. album* (Castañeda & Kendrick 1990) differ from the yellow sporodochia and filiform conidia of *L. sinense*. *Linodochium sideroxyli* is morphologically similar to *L. sinense* but differs in having wider conidia (1.5–2 μm , to 2.5 μm at the apex, 0–1 septum; Dulymamode et al. 1999) without dichotomous denticulate apices.

The significance of conidial anastomosis to form x-shaped derivative conidia in *L. sinense* is not clear. It might function as parasexual behavior or as part of sexual reproduction. However, further study is necessary to determine its function.

The teleomorph of *L. hyalinum* was found to be *Pseudohelotium pineti* (Batsch) Fuckel (Minter & Holubová-Jechová 1981). However, we found no teleomorph for *L. sinense*.

There are no DNA sequence data for any *Linodochium* species deposited in GenBank, and we were unable to isolate DNA from the *L. sinense* type specimen.

Key to species of *Linodochium*

- 1. Sporodochia yellow, amber, or orange 2
- 1. Sporodochia white to grayish white 4
- 2. Conidia 2–4-septate, $36-89 \times 1-1.5 \mu\text{m}$ *L. hyalinum*
- 2. Conidia 0–1-septate 3

3. Conidiogenous cells bearing 1–4 conidia, no denticles at apical end;
conidia 0–1-septate, $32\text{--}48 \times 1.5\text{--}2\text{ }\mu\text{m}$, wider toward apex *L. sideroxyli*
3. Conidiogenous cells bearing mostly 2 conidia, with dichotomous denticles at apex;
conidia 1-septate, $34\text{--}40 \times 1.1\text{--}1.5\text{ }\mu\text{m}$, conidial anastomosis forming x-shaped
derivative conidia *L. sinense*
4. Conidia straight, <12 -septate, $80\text{--}140 \times 2\text{--}3\text{ }\mu\text{m}$ *L. formosum*
4. Conidia falcate, 1–2-septate, $30\text{--}45 \times 1\text{--}2\text{ }\mu\text{m}$ *L. album*

New records for China

Candelabrum microsporum R.F. Castañeda & W.B. Kendr., Univ. Waterloo Biol. Ser.
35: 16 (1991)

TELEOMORPH: unknown.

CONIDIOPHORES reduced to conidiogenous cells. CONIDIOGENOUS CELLS monoblastic. CONIDIA subhyaline, solitary; conidial body multi-branched dichotomous or sometimes trichotomously, with minute apical tubercles, forming globose, subglobose or elongated conidia, $35\text{--}55\text{ }\mu\text{m}$ diam.

SPECIMEN EXAMINED: CHINA, HUBEI, Wufeng, Houhe National Nature Reserve, dead leaf of *Sycopsis sinensis*, 28 August 2008, De-Wei Li (BPI 892532).

ECOLOGY & DISTRIBUTION: On dead leaves of *Sycopsis sinensis*; *Palmae*. Previously known from Cuba; new record from China.

NOTE: Some conidia from the Chinese specimen are elongated and deviate slightly from the conidial shape of the holotype. Zhao et al. (2007) reported two other *Candelabrum* species in China, *C. brochiatum* Tubaki and *C. spinulosum* Beverw. Both species were collected on decaying branches of an undetermined tree submerged in water in Dashuwang, Mount Tianmu, Zhejiang.

Chalara affinis Sacc. & Berl., Atti Ins. Veneto Sci. Lett., Arti, Ser. 6 3: 741 (1885)

TELEOMORPH: unknown.

CONIDIOPHORES simple, erect, straight or slightly flexuous, cylindrical with an inflated base, <4 septate, sometimes constricted at the septa, pale brown, smooth, $40\text{--}70\text{ }\mu\text{m}$ long, $4\text{--}6\text{ }\mu\text{m}$ wide at the base. PHIALIDES lageniform, pale brown, smooth, $35\text{--}50 \times 4\text{--}6.5\text{ }\mu\text{m}$. CONIDIA 1-celled, colorless, smooth, cylindrical, ends flat or slightly rounded, $8\text{--}11 \times 1.5\text{--}2\text{ }\mu\text{m}$, catenate.

SPECIMEN EXAMINED: CHINA, HUBEI, Wufeng, Houhe National Nature Reserve, dead leaf of *Sycopsis sinensis*, 28 August 2008, De-Wei Li (BPI 892533).

ECOLOGY & DISTRIBUTION: On dead leaves of *Sycopsis sinensis*; *Pinus sylvestris* (Minter & Holubová-Jechová 1981). Previously known from Czech Republic and the United Kingdom; new record from China.

NOTE: the conidia from the Chinese material are slightly smaller than those of the type.

Cylindrocladium madagascariense Crous, Tax. Path. *Cylindrocladium*: 112 (2002)

TELEOMORPH: *Calonectria madagascariensis* Crous.

CONIDIOPHORES colorless, smooth, penicillate, $60\text{--}85 \times 5\text{--}6.5 \mu\text{m}$, with a septate extension, $130\text{--}350 \mu\text{m}$ long, terminating in a clavate vesicle. PHIALIDES, colorless, smooth, allantoid, cylindrical, or doliiform, $9\text{--}11 \times 3\text{--}4 \mu\text{m}$. CONIDIA cylindrical, rounded at both ends, straight, colorless, smooth, thin-walled, 3-septate, $50\text{--}55 \times 4\text{--}5 \mu\text{m}$.

SPECIMEN EXAMINED: CHINA, HUBEI, Wufeng, Houhe National Nature Reserve, dead leaf of *Sycopsis sinensis*, 20 May 2009, Yixun Wang (BPI 892534).

ECOLOGY & DISTRIBUTION: On fallen leaves of *Sycopsis sinensis*; *Eucalyptus* sp., soil. Previously known from Congo and Madagascar (Crous 2002); new record from China.

NOTES: The Chinese collection had shorter conidiophore extensions and slightly shorter conidia than those of the holotype. Crous (2002) reported that *Cylindrocladium madagascariense* is pathogenic to *Eucalyptus* sp., causing a leaf spot. It remains unknown whether this species does the same to *S. sinensis*. Further observation of *S. sinensis* foliage prior to leaf abscission is necessary to clarify this.

Subulispora britannica B. Sutton, Trans. Br. Mycol. Soc. 61(3): 422 (1973)

TELEOMORPH: unknown.

CONIDIOPHORES $20\text{--}40 \times 3.5\text{--}4 \mu\text{m}$, differentiate, solitary, straight or flexuous, 1–4 septate, smooth, thick-walled, pale brown to brown, sometimes with an enlarged base. CONIDIOGENOUS CELLS integrated, terminal polyblastic, sympodial, subhyaline to pale brown, thin-walled with 1–3 conspicuous flat unthickened scars. CONIDIA solitary, 3–10-septate, colorless, filiform, straight, thin-walled, smooth, $50\text{--}90 \mu\text{m}$ long and $2.5\text{--}3 \mu\text{m}$ wide at the base, attenuated toward the apex.

SPECIMEN EXAMINED: CHINA, HUBEI, Wufeng, Houhe National Nature Reserve, dead leaf of *Sycopsis sinensis*, 28 August 2008, De-Wei Li (BPI 892535).

ECOLOGY & DISTRIBUTION: on dead leaves of *Sycopsis sinensis*; *Hedera helix*, *Ilex aquifolium*, *I. pernyi*, and *Quercus suber* (Kirk 1982). Previously known from Russia and UK; new record from China.

NOTES: The Chinese conidiophores and conidia are slightly longer than those of the type.

New record for Mainland China

Menisporopsis novae-zelandiae S. Hughes & W.B. Kendr., N.Z. J. Bot. 6: 369 (1968)

TELEOMORPH: unknown.

SETAE solitary, dark brown, pale brown toward the apex, <17 septate, $250\text{--}400 \mu\text{m}$ long, $8\text{--}14 \mu\text{m}$ at the base. CONIDIOPHORES $40\text{--}75 \times 1.5\text{--}2 \mu\text{m}$.

CONIDIOGENOUS CELLS monophialidic, smooth, colorless or subhyaline, straight or slightly curved, $15\text{--}20 \times 3\text{--}4 \mu\text{m}$. COLLARETTE conspicuous, infundibuliform or cupulate, $2\text{--}3 \mu\text{m}$ wide, $1.5\text{--}3.5 \mu\text{m}$ deep. CONIDIA 1-septate, allantoid, colorless, smooth, $16\text{--}20 \times 2.5\text{--}3 \mu\text{m}$ with a simple setula at each end, $4\text{--}6 \mu\text{m}$ long.

SPECIMEN EXAMINED: CHINA, HUBEI, Wufeng, Houhe National Nature Reserve, dead leaf of *Sycopsis sinensis*, 20 August 2009, Yixun Wang (BPI 892536).

ECOLOGY & DISTRIBUTION: on dead leaves of *Sycopsis sinensis*; *Beilschmiedia tarairi*, *Cinnamomum osmophloeum*, *C. pedunculatum*, *C. zeylanicum*, *Cryptocarya mackinnoniana*, *Knightia excelsa*, *Litsea japonica*, palm, *Pasania edulis*, *P. kawakamii* (Hughes & Kendrick 1968, Matsushima 1980, 1989, 1993). Previously known from Australia, Ecuador, Japan, New Zealand (Matsushima 1975, 1980, 1989, 1993). This is the first report from Mainland China. Matsushima (1980) reported this species from Taiwan.

NOTES: The Chinese collection had shorter setae than those of the type and its conidiogenous cells are not polyphialidic.

Acknowledgments

The authors express their sincere gratitude to Dr. Bryce Kendrick and Dr. Rafael F. Castañeda Ruiz for their critical review of the manuscript and to Drs. Louis Magnarelli and James A. LaMondia for their pre-submission review. The authors are also grateful to Dengkui Tang, Zhizhang Zheng, Yeqing Wang, Sanshan Cai, and Jihong Yang for their assistance during the field trips. Dr. Lorelei L. Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated. This paper is dedicated to Dr. Louis Magnarelli, director of The Connecticut Agricultural Experiment Station, who passed away on July 11, 2013. DWL expresses his heartfelt gratitude to Dr. Magnarelli for his support in the past nine years.

Literature cited

- Carmichael JW. 1955. Lacto-fuchsin: a new medium for mounting fungi. *Mycologia* 47: 611.
- Castañeda RF, Kendrick B. 1990. Conidial fungi from Cuba: I. University of Waterloo, Biology Series 32: 1–53.
- Crous PW. 2002. Taxonomy and pathology of *Cylindrocladium* (*Calonectria*) and allied genera. APS Press, St. Paul MN. 278 p.
- Dang HS, Jiang MX, Tian YQ, Huang HD, Jin X. 2004. Population structure and distribution pattern of rare plant communities in Houhe Nature Reserve. *Chinese J. Appl. Ecol.* 15(12): 2206–2210.
- Dulymamode R, Kirk PM, Peeraly A. 1999. Fungi from Mauritius: three new hyphomycete species on endemic plants. *Mycotaxon* 73: 313–324.
- Dyko BJ, Sutton BC. 1979. A revision of *Linodochium*, *Pseudocenangium*, *Septopatella* and *Siroscyphella*. *Canadian Journal of Botany* 57(4): 370–385. <http://dx.doi.org/10.1139/b79-049>
- Höhnelt F von. 1909. Fragmente zur Mykologie (VIII. Mitteilung, Nr. 354 bis 406). *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften, Math.-naturw. Klasse, Abt. I*, 118: 1157–1246.
- Hughes SJ, Kendrick WB. 1968. New Zealand fungi. 12. *Menispora*, *Codinaea*, *Menisporopsis*. *New Zealand Journal of Botany* 6(3): 323–375. <http://dx.doi.org/10.1080/0028825X.1968.10428818>

- Kirk PM. 1982. New or interesting microfungi: V. Microfungi colonizing *Laurus nobilis* leaf litter. Transactions of the British Mycological Society 78(2): 293–303.
[http://dx.doi.org/10.1016/S0007-1536\(82\)80013-2](http://dx.doi.org/10.1016/S0007-1536(82)80013-2)
- Li X, Hu LL, Huang HD, Jiang MX. 2003. Structures and spatial distribution patterns of dominant populations in *Sycopsis sinensis* community in Houhe Nature Reserve. J. Appl. Ecol. 14(6): 849–852.
- Li ZZ, Huang HW, Tang DK, Wang LJ, Pu YH, Zheng CY. 2005. Situation and strategy of biodiversity conservation in the Houhe National Nature Reserve, Hubei Province, China I. Situation and study of biodiversity. Journal of Wuhan Botanical Research 23: 592–600.
- Li J, Wei YL, Dai YC. 2007. Polypores from Houhe Nature Reserve in Hubei Province. Journal of Fungal Research 5(4): 187–189.
- Li DW, Chen J, Wang Y. 2010. Two new species of dematiaceous hyphomycetes from Hubei, China. Sydowia 62 (1): 171–179.
- Li DW, Chen J, Wang Y. 2011. *Rhexodenticula zhengii* sp. nov. from fallen leaves from China. Mycotaxon 117(1): 287–290. <http://dx.doi.org/10.5248/117.287>
- Liu SX, Xu KY, Yang F-S. 1999. The biggest community *Sycopsis sinensis* was found in Houhe Nature Reserve, Hubei Province, China. Journal Huazhong Normal University (Natural Science) 33(4): 588–589.
- Matsushima T. 1975. Icones microfungorum a Matsushima lectorum. Kobe, Japan. 209 p, 415 pl.
- Matsushima T. 1980. Matsushima mycological memoirs no. 1. Kobe, Japan. 82 p..
- Matsushima T. 1989. Matsushima mycological memoirs no. 6. Kobe, Japan. 44 p.
- Matsushima T. 1993. Matsushima mycological memoirs no. 7. Kobe, Japan. 75 p.
- Minter DW, Holubová-Jechová V. 1981. New or interesting hyphomycetes on decaying pine litter from Czechoslovakia. Folia Geobotanica & Phytotaxonomica, 16 (2): 195–217.
- Song CS, Liu SX. 1999. Papers from the scientific expedition to Houhe National Nature Reserve in Hubei. Forestry Press, Beijing, China. pp. 14–33.
- Thiers BM. 2013 [continuously updated]. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> (Accessioned: 12.02.2013)
- Wu WP, Zhuang WY. 2005. *Sporidesmium*, *Endophragmiella* and related genera from China, Fungal Diversity Research Series 15. 351 p.
- Zhao G, Liu X, Wu W. 2007. Helicosporous hyphomycetes from China. Fungal Diversity 26: 313–524.
- Zhuang, WY, Y Nong, J Luo. 2007. New species and new Chinese records of *Bionectriaceae* and *Nectriaceae* (*Hypocreales*, *Ascomycetes*) from Hubei, China. Fungal Diversity 24: 347–357.